


Sobolev Spaces, Integral Equations, and Scattering on non-Lipschitz and Fractal Sets

Simon Chandler-Wilde,

University of Reading



Part 3 Sobolev Spaces

Tempered Distributions

$$C_0^\infty(\mathbb{R}^n) \subset \mathcal{S}(\mathbb{R}^n) \subset C^\infty(\mathbb{R}^n)$$

space of rapidly
decreasing C^∞ fns

Tempered Distributions

$$C_0^\infty(\mathbb{R}^n) \subset \mathcal{S}(\mathbb{R}^n) \subset C^\infty(\mathbb{R}^n)$$

space of rapidly
decreasing C^∞ fns

$\mathcal{S}'(\mathbb{R}^n)$, dual space of $\mathcal{S}(\mathbb{R}^n)$ - "tempered
distributions"

$\langle u, v \rangle \in \mathbb{C}$, for $u \in \mathcal{S}'(\mathbb{R}^n)$, $v \in \mathcal{S}(\mathbb{R}^n)$
is action of linear functional u on v

Tempered Distributions

$$C_0^\infty(\mathbb{R}^n) \subset \mathcal{S}(\mathbb{R}^n) \subset C^\infty(\mathbb{R}^n)$$

space of rapidly decreasing C^∞ fns

$\mathcal{S}'(\mathbb{R}^n)$, dual space of $\mathcal{S}(\mathbb{R}^n)$ - "tempered distributions"

$$\langle u, v \rangle \in \mathbb{C}, \text{ for } u \in \mathcal{S}'(\mathbb{R}^n), v \in \mathcal{S}(\mathbb{R}^n)$$

is action of linear functional u on v

$\hat{u} = \mathcal{F}u$ is FT of u , \mathcal{F} an isomorphism on $\mathcal{S}(\mathbb{R}^n)$ and on $\mathcal{S}'(\mathbb{R}^n)$

$$\langle \hat{u}, v \rangle := \langle u, \hat{v} \rangle, u \in \mathcal{S}', v \in \mathcal{S}$$

Ex Given $z \in \mathbb{R}^n$ define

$\delta_z \in \mathcal{S}'(\mathbb{R}^n)$ by

$$\langle \delta_z, v \rangle = v(z), \quad v \in \mathcal{S}(\mathbb{R}^n)$$

Show that $\mathcal{F} \delta_z = f$, where

$$f(\xi) = (2\pi)^{-n/2} e^{-i z \cdot \xi}, \quad \xi \in \mathbb{R}^n$$

$$\mathcal{F} u(\xi) = \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{-i \xi \cdot x} u(x) dx, \quad \xi \in \mathbb{R}^n, \quad u \in \mathcal{S}(\mathbb{R}^n)$$

Sobolev spaces on \mathbb{R}^n

For $s \geq 0$, $H^s(\mathbb{R}^n) = \{u \in L^2(\mathbb{R}^n) \mid \|u\|_{H^s} < \infty\}$

where

$$\|u\|_{H^s}^2 = \int_{\mathbb{R}^n} (1 + |\xi|^2)^s |\hat{u}(\xi)|^2 d\xi$$

Sobolev spaces on \mathbb{R}^n

For $s \geq 0$, $H^s(\mathbb{R}^n) = \{v \in L^2(\mathbb{R}^n) \mid \|v\|_{H^s} < \infty\}$

where

$$\|v\|_{H^s}^2 = \int_{\mathbb{R}^n} (1 + |\xi|^2)^s |\hat{v}(\xi)|^2 d\xi$$

$$\|v\|_{H^0}^2 = \int_{\mathbb{R}^n} |\hat{v}(\xi)|^2 d\xi = \int_{\mathbb{R}^n} |v(x)|^2 dx = \|v\|_{L^2}^2$$

$$\text{so } H^0(\mathbb{R}^n) = L^2(\mathbb{R}^n)$$

Sobolev spaces on \mathbb{R}^n

For $s \geq 0$, $H^s(\mathbb{R}^n) = \{u \in L^2(\mathbb{R}^n) \mid \|u\|_{H^s} < \infty\}$

where

$$\|u\|_{H^s}^2 = \int_{\mathbb{R}^n} (1 + |\xi|^2)^s |\hat{u}(\xi)|^2 d\xi$$

$$\begin{aligned} \widehat{\partial_j u}(\xi) &= i \xi_j \hat{u}(\xi) \Rightarrow \|u\|_{H^1}^2 = \int_{\mathbb{R}^n} (1 + \xi_1^2 + \dots + \xi_n^2) |\hat{u}(\xi)|^2 d\xi \\ &= \int_{\mathbb{R}^n} (|\hat{u}|^2 + \sum_j |\widehat{\partial_j u}|^2) \\ &= \int_{\mathbb{R}^n} (|u|^2 + |\nabla u|^2) \end{aligned}$$

Sobolev spaces on \mathbb{R}^n

For $s \geq 0$, $H^s(\mathbb{R}^n) = \{u \in L^2(\mathbb{R}^n) \mid \|u\|_{H^s} < \infty\}$

where

$$\|u\|_{H^s}^2 = \int_{\mathbb{R}^n} (1 + |\xi|^2)^s |\hat{u}(\xi)|^2 d\xi$$

For $s > 0$, $H^{-s}(\mathbb{R}^n) = (H^s(\mathbb{R}^n))^*$

Sobolev spaces on \mathbb{R}^n

For $s \geq 0$, $H^s(\mathbb{R}^n) = \{u \in L^2(\mathbb{R}^n) \mid \|u\|_{H^s} < \infty\}$

where

$$\|u\|_{H^s}^2 = \int_{\mathbb{R}^n} (1 + |\xi|^2)^s |\hat{u}(\xi)|^2 d\xi$$

For $s > 0$, $H^{-s}(\mathbb{R}^n) = (H^s(\mathbb{R}^n))^*$

For $s > 0$, $S(\mathbb{R}^n) \subset H^s(\mathbb{R}^n)$, so $H^{-s}(\mathbb{R}^n) \subset S^*(\mathbb{R}^n)$

Sobolev spaces on \mathbb{R}^n

For $s \geq 0$, $H^s(\mathbb{R}^n) = \{u \in L^2(\mathbb{R}^n) \mid \|u\|_{H^s} < \infty\}$

where

$$\|u\|_{H^s}^2 = \int_{\mathbb{R}^n} (1 + |\xi|^2)^s |\hat{u}(\xi)|^2 d\xi$$

For $s > 0$, $H^{-s}(\mathbb{R}^n) = (H^s(\mathbb{R}^n))^*$

For $s > 0$, $\mathcal{S}(\mathbb{R}^n) \subset H^s(\mathbb{R}^n)$, so $H^{-s}(\mathbb{R}^n) \subset \mathcal{S}'(\mathbb{R}^n)$,
indeed

$$H^{-s}(\mathbb{R}^n) = \{u \in \mathcal{S}'(\mathbb{R}^n) \mid \|u\|_{H^{-s}} < \infty\}$$

↑ Proof just uses Riesz rep thm on $L^2(\mathbb{R}^n)$

Sobolev spaces on \mathbb{R}^n

For $s \geq 0$, $H^s(\mathbb{R}^n) = \{u \in L^2(\mathbb{R}^n) \mid \|u\|_{H^s} < \infty\}$

where

$$\|u\|_{H^s}^2 = \int_{\mathbb{R}^n} (1 + |\xi|^2)^s |\hat{u}(\xi)|^2 d\xi$$

For $s > 0$, $H^{-s}(\mathbb{R}^n) = (H^s(\mathbb{R}^n))^*$

$$= \{u \in \mathcal{S}'(\mathbb{R}^n) \mid \|u\|_{H^{-s}} < \infty\}$$

Ex Show $\delta_z \in H^{-s}(\mathbb{R}^n)$

$$\iff s > n/2$$

Closed subspaces of $H^s(\mathbb{R}^n)$

For open $\Omega \subset \mathbb{R}^n$,

$$\tilde{H}^s(\Omega) = \overline{C_0^\infty(\Omega)}^{H^s(\mathbb{R}^n)}$$

Closed subspaces of $H^s(\mathbb{R}^n)$

For **open** $\Omega \subset \mathbb{R}^n$,

$$\tilde{H}^s(\Omega) = \overline{C_0^\infty(\Omega)}^{H^s(\mathbb{R}^n)}$$

For **closed** $F \subset \mathbb{R}^n$,

$$H_F^s := \{v \in H^s(\mathbb{R}^n) \cdot \text{supp}(v) \subset F\}$$

Closed subspaces of $H^s(\mathbb{R}^n)$

For **open** $\Omega \subset \mathbb{R}^n$,

$$\tilde{H}^s(\Omega) = \overline{C_0^\infty(\Omega)}^{H^s(\mathbb{R}^n)}$$

For **closed** $F \subset \mathbb{R}^n$,

$$H_F^s := \{u \in H^s(\mathbb{R}^n) \cdot \text{supp}(u) \subset F\}$$

NB $u \in H_F^s \Leftrightarrow \langle u, v \rangle = 0, \forall v \in C_0^\infty(\mathbb{R}^n \setminus F)$

Closed subspaces of $H^s(\mathbb{R}^n)$

For **open** $\Omega \subset \mathbb{R}^n$,

$$\tilde{H}^s(\Omega) = \overline{C_0^\infty(\Omega)}^{H^s(\mathbb{R}^n)}$$

For **closed** $F \subset \mathbb{R}^n$,

$$H_F^s := \{v \in H^s(\mathbb{R}^n) \cdot \text{supp}(v) \subset F\}$$

$$\tilde{H}^s(\Omega) \subset H_{\Omega}^s$$

Closed subspaces of $H^s(\mathbb{R}^n)$

For **open** $\Omega \subset \mathbb{R}^n$

$$\tilde{H}^s(\Omega) = \overline{C_0^\infty(\Omega)}^{H^s(\mathbb{R}^n)}$$



For **closed** $F \subset \mathbb{R}^n$,

$$H_F^s := \{v \in H^s(\mathbb{R}^n) : \text{supp}(v) \subset F\}$$

$$\tilde{H}^s(\Omega) \subset H_{\Omega}^s \quad - \quad \text{often } \tilde{H}^s(\Omega) = H_{\Omega}^s$$




Thm $\tilde{H}^s(\Omega) = H^s_{\mathbb{R}} \quad \text{if}$

(1) Ω is C^0
and $s \in \mathbb{R}$




Thm $\tilde{H}^s(\Omega) = H^s_\Omega$ if

(ii) Ω is C^0 except at a finite number of points on $\partial\Omega$ and $|s| \leq \frac{1}{2}$ ($n=1$), $|s| \leq 1$ ($n \geq 2$)



Thm $\tilde{H}^s(\Omega) = H^s_\Omega$ if

(iii) Ω is C^0 except at a countable set of points on $\partial\Omega$ with finitely many limit points and $|S| \leq \frac{1}{2}$ ($n=1$), $|S| \leq 1$ ($n \geq 2$)




(C-w, Hewitt,
Moiola,
IEOT 2017)

Thm $\tilde{H}^s(\Omega) = H^s_\Omega$ if

(iii) Ω is in a specific set of domains

with fractal boundary,

(Caetano, Hevett, Moiola 2019) $s \in \mathbb{R}$



Thm If $\Omega \subset \mathbb{R}^n$ open
and $\tilde{\Omega} = \mathbb{R}^n$ (so $H_{\mathbb{R}^n}^s = H^s(\mathbb{R}^n)$)
then

$$\tilde{H}^s(\Omega) = H_{\mathbb{R}^n}^s \iff H_{\partial\Omega}^{-s} = \{0\}$$

Thm If $\Omega \subset \mathbb{R}^n$ open
and $\tilde{\Omega} = \mathbb{R}^n$ (so $H^s_{\tilde{\Omega}} = H^s(\mathbb{R}^n)$)
then

$$\tilde{H}^s(\Omega) = H^s_{\tilde{\Omega}} \iff H^{-s}_{\partial\Omega} = \{0\}$$

Ex If $z \in \mathbb{R}^n$ and $\Omega = \mathbb{R}^n \setminus \{z\}$,
so $\partial\Omega = \{z\}$,

$$\tilde{H}^s(\Omega) = H^s_{\tilde{\Omega}} \iff H^{-s}_{\partial\Omega} = \{0\} \iff s \leq \frac{n}{2}$$

Thm If $\Omega \subset \mathbb{R}^n$ open
and $\tilde{\Omega} = \mathbb{R}^n$ (so $H^s_{\tilde{\Omega}} = H^s(\mathbb{R}^n)$)
then

$$\tilde{H}^s(\Omega) = H^s_{\tilde{\Omega}} \iff H^{-s}_{\partial\Omega} = \{0\}$$

Proof \Rightarrow Suppose $\phi \in H^{-s}_{\partial\Omega}$ &

$\tilde{H}^s(\Omega) = H^s(\mathbb{R}^n)$ Then

$$\forall \psi \in H^s(\mathbb{R}^n), \langle \phi, \psi \rangle = \lim_{n \rightarrow \infty} \langle \phi, \psi_n \rangle = 0$$

$\psi_n \in C^\infty_0(\Omega)$

Thm If $\Omega \subset \mathbb{R}^n$ open
and $\tilde{\Omega} = \mathbb{R}^n$ (so $H_{\tilde{\Omega}}^s = H^s(\mathbb{R}^n)$)
then

$$\tilde{H}^s(\Omega) = H_{\tilde{\Omega}}^s \iff H_{\partial\Omega}^{-s} = \{0\}$$

Proof \Leftarrow Exercise!

(Maz'ya, "Sobolev Spaces with",
Thm 13.2.1)

Thm If $\Omega \subset \mathbb{R}^n$ open
and $\tilde{\Omega} = \mathbb{R}^n$ (so $H_{\mathbb{R}}^s = H^s(\mathbb{R}^n)$)
then

$$\tilde{H}^s(\Omega) = H_{\mathbb{R}}^s \iff H_{\partial\Omega}^{-s} = \{0\}$$

$$\implies \dim_{\mathbb{H}} \partial\Omega \leq n - 2s$$

($\frac{n}{2} \geq s > 0$)

Hausdorff dim $\in [0, n]$

Open problem

Necessary and sufficient conditions
for $H^s(\Omega) = H^s_{\bar{\Omega}}$
for general open $\Omega \subset \mathbb{R}^n$, $s \in \mathbb{R}$

(Review of what known in
C-W, Hewett, Moiola, IEOT, 2017)

Open problem

True \forall
Jones domains?

Necessary and sufficient conditions
for


$$H^s(\Omega) = H^s_{\bar{\Omega}}$$

for general open $\Omega \subset \mathbb{R}^n$, $s \in \mathbb{R}$

(Review of what known in
C-W, Hewett, Moiola, IEOT, 2017)

Sequences of subspaces and Mosco convergence

Suppose
 $\Omega_1 \subset \Omega_2 \subset$
are open and let
 $V_j := \tilde{H}^s(\Omega_j)$ so
 $V_1 \subset V_2 \subset$




Sequences of subspaces and Mosco convergence

Suppose
 $\Omega_1 \subset \Omega_2 \subset$
are open and let

$V_j := \tilde{H}^s(\Omega_j)$ so

$V_1 \subset V_2 \subset$

Then
 $V_j \xrightarrow{M} V := \bigcup_j V_j = \bigcup_j \tilde{H}^s(\Omega_j)$




Sequences of subspaces and Mosco convergence

Suppose
 $\Omega_1 \subset \Omega_2 \subset \dots$
are open and let
 $V_j := \tilde{H}^s(\Omega_j)$ so

$$V_1 \subset V_2 \subset \dots$$

Then

$$V_j \xrightarrow{M} V := \bigcup_j V_j = \bigcup_j \tilde{H}^s(\Omega_j)$$



Ex show

$$V = \tilde{H}^s\left(\bigcup_j \Omega_j\right)$$

Sequences of subspaces and Mosco convergence


Suppose

$F_1 \supset F_2 \supset \dots$
are closed and let

$V_j := H_{F_j}^s$ so

$V_1 \supset V_2 \supset \dots$ Then

$V_j \xrightarrow{M} V := \bigcap_j V_j = \bigcap_j H_{F_j}^s = H_F^s, F := \bigcap_j F_j$



Sequences of subspaces and Mosco convergence

Suppose

$$\Omega_1 \supset \Omega_2 \supset$$


are open and let

$$V_j := \tilde{H}^s(\Omega_j) \text{ so}$$

$$V_1 \supset V_2 \supset$$

Then

$$V_j \xrightarrow{M} V := \bigcap_j V_j = \bigcap_j \tilde{H}^s(\Omega_j) = ??$$



Part 4: Integral Equation
Formulations for Scattering
by Thin Screens*

* which may be fractal!

Given $g \in H^{-1}(\mathbb{R}^2) := (H^1(\mathbb{R}^2))^*$
find $v \in \tilde{H}^1(D) = \frac{C_0^\infty(D)}{H^1(\mathbb{R}^2)}$
st $\Delta v + k^2 v = g$ in D

$k \in \mathbb{C}$,
 $\text{Im} k > 0$

\mathcal{X}_2

$D := \mathbb{R}^2 \setminus \Gamma$

Γ , closed \mathcal{X}_1

Given $g \in H^{-1}(\mathbb{R}^2) := (H^1(\mathbb{R}^2))^*$
find $v \in \tilde{H}^1(D) = \frac{C_0^\infty(D)}{H^1(\mathbb{R}^2)}$
st $\Delta u + k^2 u = g$ in D

\Leftrightarrow Find $v \in \tilde{H}^1(D)$ st $a(u, v) = \langle g, \bar{v} \rangle, \forall v \in \tilde{H}^1(D)$

Given $g \in H^{-1}(\mathbb{R}^2) := (H^1(\mathbb{R}^2))^*$
find $v \in \tilde{H}^1(D) = \frac{C_0^\infty(D)}{H^1(\mathbb{R}^2)}$
st $\Delta u + k^2 u = g$ in D

\Leftrightarrow Find $v \in \tilde{H}^1(D)$ st $a(u, v) = \langle g, \bar{v} \rangle, \forall v \in \tilde{H}^1(D)$

Applying L-M these have exactly one soln
and $\|u\| \leq c(k) \|g\|$

Given $g \in H^{-1}(\mathbb{R}^2) := (H^1(\mathbb{R}^2))^*$
find $v \in \tilde{H}^1(D) = \frac{C_0^\infty(D)}{H^1(\mathbb{R}^2)}$
st $\Delta u + k^2 u = g$ in D

\Leftrightarrow Find $v \in \tilde{H}^1(D)$ st $a(u, v) = \langle g, \bar{v} \rangle, \forall v \in \tilde{H}^1(D)$

Applying L-M these have exactly one soln
and $\|u\| \leq c(k) \|g\|$ TRUE IF $D = \mathbb{R}^2$

Given $g \in H^{-1}(\mathbb{R}^2) := (H^1(\mathbb{R}^2))^*$,
find $v \in H^1(\mathbb{R}^2)$
st $\Delta u + k^2 u = g$ in \mathbb{R}^2

\Leftrightarrow Find $v \in H^1(\mathbb{R}^2)$ st $a(u, v) = \langle g, \tilde{v} \rangle, \forall v \in H^1(\mathbb{R}^2)$

Applying L-M these have exactly one soln
and $\|u\| \leq c(k) \|g\|$ TRUE IF $D = \mathbb{R}^2$

Given $g \in H^{-1}(\mathbb{R}^2) := (H^1(\mathbb{R}^2))^*$,
find $v \in H^1(\mathbb{R}^2)$
st $\Delta v + k^2 v = g$ in \mathbb{R}^2

\Leftrightarrow Find $v \in H^1(\mathbb{R}^2)$ st $a(u, v) = \langle g, \bar{v} \rangle, \forall v \in H^1(\mathbb{R}^2)$

Applying L-M these have exactly one soln
and $\|v\| \leq c(k) \|g\|$ So $\Delta + k^2: H^1(\mathbb{R}^2) \rightarrow H^{-1}(\mathbb{R}^2)$
is isomorphism with inverse $G = (\Delta + k^2)^{-1}$ & $\|G\| \leq c(k)$

Given $g \in H^{-1}(\mathbb{R}^2) := (H^1(\mathbb{R}^2))^*$,

find $v \in H^1(\mathbb{R}^2)$

st

$$\Delta u + k^2 u = g \iff u = Gg$$

\iff Find $v \in H^1(\mathbb{R}^2)$ st $a(u, v) = \langle g, \bar{v} \rangle, \forall v \in H^1(\mathbb{R}^2)$

Applying L-M these have exactly one soln

and $\|u\| \leq c(k) \|g\|$ So $\Delta + k^2 : H^1(\mathbb{R}^2) \rightarrow H^{-1}(\mathbb{R}^2)$

is isomorphism with inverse $G = (\Delta + k^2)^{-1}$ & $\|G\| \leq c(k)$

$$\begin{aligned} &\text{Given } g \in H^{-1}(\mathbb{R}^2) := (H^1(\mathbb{R}^2))^* \\ &\text{find } v \in \tilde{H}^1(D) = \frac{C_0^\infty(D)}{H^1(\mathbb{R}^2)} \\ &\text{st } \Delta v + k^2 v = g \text{ in } D \end{aligned}$$

Let $\phi = \Delta v + k^2 v - g$ Then $\phi = 0$ in D , so
 $\phi \in H_0^{-1}$

$$\begin{aligned} &\text{Given } g \in H^{-1}(\mathbb{R}^2) := (H^1(\mathbb{R}^2))^* \\ &\text{find } v \in \tilde{H}^1(D) = \frac{C_0^\infty(D)}{H^1(\mathbb{R}^2)} \\ &\text{st } \Delta v + k^2 v = g \text{ in } D \end{aligned}$$

Let $\phi = \Delta v + k^2 v - g$. Then $\phi = 0$ in D , so $\phi \in H_{\Gamma}^{-1}$, and

$$G \phi = G(\Delta + k^2)v - Gg = v - Gg$$

$$\begin{aligned} &\text{Given } g \in H^{-1}(\mathbb{R}^2) := (H^1(\mathbb{R}^2))^* \\ &\text{find } u \in \tilde{H}^1(D) = \frac{C_0^\infty(D)}{H^1(\mathbb{R}^2)} \\ &\text{st } \Delta u + k^2 u = g \text{ in } D \end{aligned}$$

Let $\phi = \Delta u + k^2 u - g$. Then $\phi = 0$ in D , so $\phi \in H_{\Gamma}^{-1}$, and

$$\langle \phi, v \rangle = \langle (\Delta + k^2)u - g, v \rangle = \langle u, \Delta v - g \rangle$$

Now $\langle \psi, v \rangle = 0, \forall v \in C_0^\infty(D), \psi \in H_{\Gamma}^{-1}$

$$\begin{aligned} &\text{Given } g \in H^{-1}(\mathbb{R}^2) := (H^1(\mathbb{R}^2))^* \\ &\text{find } v \in \tilde{H}^1(D) = \frac{C_0^\infty(D)}{H^1(\mathbb{R}^2)} \\ &\text{st } \Delta v + k^2 v = g \text{ in } D \end{aligned}$$

Let $\phi = \Delta v + k^2 v - g$. Then $\phi = 0$ in D , so $\phi \in H_{\Gamma}^{-1}$, and

$$\int \phi = \int (\Delta + k^2)v - \int g = 0 - \int g$$

Now $\langle \psi, v \rangle = 0$, $\forall v \in C_0^\infty(D)$, $\psi \in H_{\Gamma}^{-1}$
 $\tilde{H}^1(D)$

$$\begin{aligned} &\text{Given } g \in H^{-1}(\mathbb{R}^2) := (H^1(\mathbb{R}^2))^* \\ &\text{find } v \in \tilde{H}^1(D) = \frac{C_0^\infty(D)}{H^1(\mathbb{R}^2)} \\ &\text{st } \Delta v + k^2 v = g \text{ in } D \end{aligned}$$

Let $\phi = \Delta v + k^2 v - g$. Then $\phi = 0$ in D , so $\phi \in H_{\Gamma}^{-1}$, and

$$\int \phi = \int (\Delta + k^2)v - \int g = 0 - \int g$$

Now $\langle \psi, v \rangle = 0$, $\forall v \in C_0^\infty(D)$, $\psi \in H_{\Gamma}^{-1}$

So $\langle \psi, \int \phi \rangle = -\langle \psi, \int g \rangle$, $\tilde{H}^1(D)$
 $\forall \psi \in H_{\Gamma}^{-1}$

$$\begin{aligned} &\text{Given } g \in H^{-1}(\mathbb{R}^2) := (H^1(\mathbb{R}^2))^* \\ &\text{find } u \in \tilde{H}^1(D) = \overline{C_0^\infty(D)}^{H^1(\mathbb{R}^2)} \\ &\text{st } \Delta u + k^2 u = g \text{ in } D \end{aligned}$$

Let $\phi = \Delta u + k^2 u - g$. Then $\phi = 0$ in D , so $\phi \in H_{\Gamma}^{-1}$, and

$$\chi \phi = \chi(\Delta + k^2)u - \chi g = u - \chi g$$

Now $\langle \psi, v \rangle = 0$, $\forall v \in \tilde{H}^1(D)$, $\psi \in H_{\Gamma}^{-1}$

Find $\phi \in H_{\Gamma}^{-1}$ st $\langle \psi, \chi \phi \rangle = -\langle \psi, \chi g \rangle$, $\forall \psi \in H_{\Gamma}^{-1}$

$$\begin{aligned} &\text{Given } g \in H^{-1}(\mathbb{R}^2) := (H^1(\mathbb{R}^2))^* \\ &\text{find } u \in \tilde{H}^1(D) = \overline{C_0^\infty(D)}^{H^1(\mathbb{R}^2)} \\ &\text{st } \Delta u + k^2 u = g \text{ in } D \end{aligned}$$

Let $\phi = \Delta u + k^2 u - g$. Then $\phi = 0$ in D , so $\phi \in H_{\Gamma}^{-1}$, and

$$\mathcal{G}\phi = \mathcal{G}(\Delta + k^2)u - \mathcal{G}g = u - \mathcal{G}g$$

Now $\langle \psi, v \rangle = 0, \forall v \in \tilde{H}^1(D), \psi \in H_{\Gamma}^{-1}$

Find $\phi \in H_{\Gamma}^{-1}$ st $\langle \psi, \mathcal{G}\phi \rangle = -\langle \psi, \mathcal{G}g \rangle, \forall \psi \in H_{\Gamma}^{-1}$

Our INTEGRAL EQ V P

(1)

$$\begin{aligned} &\text{Given } g \in H^{-1}(\mathbb{R}^2) := (H^1(\mathbb{R}^2))^* \\ &\text{find } u \in \tilde{H}^1(D) = \overline{C_0^\infty(D)}^{H^1(\mathbb{R}^2)} \\ &\text{st } \Delta u + k^2 u = g \text{ in } D \end{aligned}$$

(2) Find $\phi \in H_{\Gamma}^{-1}$ st $\langle \psi, G\phi \rangle = -\langle \psi, Gg \rangle, \forall \psi \in H_{\Gamma}^{-1}$

Thm

If u satisfies (1) then
 $\phi = \Delta u + k^2 u - g$ satisfies (2) and
 $u = Gg + G\phi$

(1)

$$\begin{aligned} &\text{Given } g \in H^{-1}(\mathbb{R}^2) := (H^1(\mathbb{R}^2))^* \\ &\text{find } u \in \tilde{H}^1(D) = \overline{C_0^\infty(D)}^{H^1(\mathbb{R}^2)} \\ &\text{st } \Delta u + k^2 u = g \text{ in } D \end{aligned}$$

(2) Find $\phi \in H_0^{-1}$ st $\langle \psi, G\phi \rangle = -\langle \psi, Gg \rangle, \forall \psi \in H_0^{-1}$

Thm

If u satisfies (1) then

$\phi = \Delta u + k^2 u - g$ satisfies (2) and

$$u = Gg + G\phi \quad (+)$$

Conversely, if ϕ satisfies (2) then u given by (+) satisfies (1)

(1)

$$\begin{aligned} &\text{Given } g \in H^{-1}(\mathbb{R}^2) := (H^1(\mathbb{R}^2))^* \\ &\text{find } u \in \tilde{H}^1(D) = \overline{C_0^\infty(D)}^{H^1(\mathbb{R}^2)} \\ &\text{st } \Delta u + k^2 u = g \text{ in } D \end{aligned}$$

$$(2) \text{ Find } \phi \in \tilde{H}_\Gamma^{-1} \text{ st } \langle \psi, G\phi \rangle = -\langle \psi, Gg \rangle, \forall \psi \in \tilde{H}_\Gamma^{-1}$$

Thm

If u satisfies (1) then

$$\phi = \Delta u + k^2 u - g \text{ satisfies (2) and}$$

$$u = Gg + G\phi \quad (+)$$

(A) proved already

Conversely, if ϕ satisfies (2) then u given by (+) satisfies (1)

Follows from (A) and well-posedness of (2)

Well-posedness of our I E formulation

Given $g \in H^{-1}(\mathbb{R}^2)$ find $\phi \in H^{-1}_\Gamma$ s.t. $\langle \psi, \mathcal{G}\phi \rangle = -\langle \psi, \mathcal{G}g \rangle, \forall \psi \in H^{-1}_\Gamma$

\Leftrightarrow Find $\phi \in H^{-1}_\Gamma$ s.t. $A(\phi, \psi) = -\langle \bar{\psi}, \mathcal{G}g \rangle,$
where $\forall \psi \in H^{-1}_\Gamma$
 $A(\phi, \psi) = \langle \bar{\psi}, \mathcal{G}\phi \rangle \quad \phi, \psi \in H^{-1}_\Gamma$

Well-posedness of our I E formulation

Given $g \in H^{-1}(\mathbb{R}^2)$ find $\phi \in H^{-1}_\Gamma$ s.t. $\langle \psi, \mathcal{G}\phi \rangle = -\langle \psi, \mathcal{G}g \rangle, \forall \psi \in H^{-1}_\Gamma$

\Leftrightarrow Find $\phi \in H^{-1}_\Gamma$ s.t. $A(\phi, \psi) = -\langle \bar{\psi}, \mathcal{G}g \rangle,$
where $\forall \psi \in H^{-1}_\Gamma$

$$A(\phi, \psi) = \langle \bar{\psi}, \mathcal{G}\phi \rangle, \quad \phi, \psi \in H^{-1}_\Gamma$$

Continuous? Coercive?

$$A(\phi, \psi) = \langle \bar{\psi}, G\phi \rangle \quad \phi, \psi \in H_r^{-1}$$

Continuous?

$$A(\phi, \psi) = \langle \bar{\psi}, G\phi \rangle \quad \phi, \psi \in H_r^{-1}$$

Continuous?

$$\forall \phi, \psi \in H_r^{-1},$$

$$\begin{aligned} |A(\phi, \psi)| &= |\langle \bar{\psi}, G\phi \rangle| \leq \|\psi\| \|G\phi\| \\ &\leq c(k) \|\psi\| \|\phi\| \end{aligned}$$

$$A(\phi, \psi) = \langle \bar{\psi}, G\phi \rangle \quad \phi, \psi \in H_1^{-1}$$

Coercive?

Given $g \in H^{-1}(\mathbb{R}^2) := (H^1(\mathbb{R}^2))^*$,

find $v \in H^1(\mathbb{R}^2)$

st

$$\Delta u + k^2 u = g \iff u = Gg$$

\iff Find $v \in H^1(\mathbb{R}^2)$ st $a(u, v) = \langle g, v \rangle, \forall v \in H^1(\mathbb{R}^2)$

$$a(u, v) = \int_{\mathbb{R}^2} (k^2 u \bar{v} - \nabla u \cdot \nabla \bar{v})$$

Given $g \in H^{-1}(\mathbb{R}^2) := (H^1(\mathbb{R}^2))^*$,

find $v \in H^1(\mathbb{R}^2)$

st

$$\Delta u + k^2 u = g \iff u = Gg$$

\iff Find $v \in H^1(\mathbb{R}^2)$ st $a(u, v) = \langle g, \bar{v} \rangle, \forall v \in H^1(\mathbb{R}^2)$

coercive

Soln is $v = Gg$ so

$$a(Gg, Gg) = \langle g, \overline{Gg} \rangle$$

$$A(\phi, \psi) = \langle \bar{\psi}, \mathcal{G}\phi \rangle \quad \phi, \psi \in H_{\Gamma}^{-1}$$

Coercive?

$$\forall \phi \in H_{\Gamma}^{-1},$$

$$\begin{aligned} |A(\phi, \phi)| &= |\langle \bar{\phi}, \mathcal{G}\phi \rangle| \\ &= |a(\mathcal{G}\phi, \mathcal{G}\phi)| \end{aligned}$$

$$A(\phi, \psi) = \langle \bar{\psi}, \mathcal{L}\phi \rangle \quad \phi, \psi \in H_{\Gamma}^{-1}$$

Coercive?

$$\forall \phi \in H_{\Gamma}^{-1},$$

$$|A(\phi, \phi)| = |\langle \bar{\phi}, \mathcal{L}\phi \rangle|$$

Coercivity
of $a(\cdot, \cdot)$

$$= |a(\mathcal{L}\phi, \mathcal{L}\phi)|$$
$$\geq c(k) \|\mathcal{L}\phi\|^2$$

$$A(\phi, \psi) = \langle \bar{\psi}, G\phi \rangle \quad \phi, \psi \in H_r^{-1}$$

Coercive?

$$\forall \phi \in H_r^{-1},$$

since

$$\|\phi\| = \|(\Delta + k^2)G\phi\| \leq \|\Delta + k^2\| \|G\phi\|$$

$$|A(\phi, \phi)| = |\langle \bar{\phi}, G\phi \rangle|$$

$$= |a(G\phi, G\phi)|$$

$$\geq c(k) \|G\phi\|^2 \geq \tilde{c}(k) \|\phi\|^2$$

Well-posedness of our I E formulation

Given $g \in H^1(\mathbb{R}^2)$ find $\phi \in H_\Gamma^{-1}$ s.t. $\langle \psi, \mathcal{G}\phi \rangle = -\langle \psi, \mathcal{G}g \rangle, \forall \psi \in H_\Gamma^{-1}$

\Leftrightarrow Find $\phi \in H_\Gamma^{-1}$ s.t. $A(\phi, \psi) = -\langle \bar{\psi}, \mathcal{G}g \rangle,$
where $\forall \psi \in H_\Gamma^{-1}$

$$\underline{A(\phi, \psi)} = \langle \bar{\psi}, \mathcal{G}\phi \rangle \quad \phi, \psi \in H_\Gamma^{-1}$$

Cont γ coercive so

Well-posedness of our I E formulation

Given $g \in H^1(\mathbb{R}^2)$ find $\phi \in H_\Gamma^{-1}$ s.t. $\langle \psi, \mathcal{L}\phi \rangle = -\langle \psi, \mathcal{L}g \rangle, \forall \psi \in H_\Gamma^{-1}$

\Leftrightarrow Find $\phi \in H_\Gamma^{-1}$ s.t. $A(\phi, \psi) = -\langle \bar{\psi}, \mathcal{L}g \rangle,$
where $\forall \psi \in H_\Gamma^{-1}$

$$A(\phi, \psi) = \langle \bar{\psi}, \mathcal{L}\phi \rangle, \quad \phi, \psi \in H_\Gamma^{-1}$$

Thm \exists exactly one soln $\phi \in H_\Gamma^{-1}$ and

$$\|\phi\| \leq c(\kappa) \|\mathcal{L}g\| \leq \tilde{c}(\kappa) \|g\|$$

Proof L-M

Well-posedness of our I E formulation

Given $g \in H^1(\mathbb{R}^2)$ find $\phi \in H_\Gamma^{-1}$ s.t. $\langle \psi, \mathcal{G}\phi \rangle = -\langle \psi, \mathcal{G}g \rangle, \forall \psi \in H_\Gamma^{-1}$

Well-posedness of our I E formulation

Given $g \in \tilde{H}^{-1}(\mathbb{R}^2)$ find $\phi \in \tilde{H}^{-1}_\Gamma$ st $\langle \psi, \mathcal{G}\phi \rangle = -\langle \psi, \mathcal{G}g \rangle, \forall \psi \in \tilde{H}^{-1}_\Gamma$

OK . but where is
the \int integral ?
boundary

Thm (e.g. McLean, 2000, p104)

$$H_{\Gamma}^{-1}(\mathbb{R}^2) = \{\gamma^* \phi \cdot \phi \in H_{\Gamma}^{-1/2}(\mathbb{R})\}$$

where $\gamma^* : H^{-1/2}(\mathbb{R}) \rightarrow H^{-1}(\mathbb{R}^2)$,

defined by

$$\langle \gamma^* \phi, \psi \rangle := \langle \phi, \gamma \psi \rangle, \quad \phi \in H^{-1/2}(\mathbb{R}), \psi \in H^1(\mathbb{R}^2)$$

Thm (e.g. McLean, 2000, p104)

$$H_{\Gamma}^{-1}(\mathbb{R}^2) = \{\gamma^* \phi \cdot \phi \in H_{\Gamma}^{-1/2}(\mathbb{R})\}$$

where $\gamma^* : H^{-1/2}(\mathbb{R}) \rightarrow H^{-1}(\mathbb{R}^2)$,

defined by

$$\langle \gamma^* \phi, \psi \rangle := \langle \phi, \gamma \psi \rangle, \quad \phi \in H^{-1/2}(\mathbb{R}), \psi \in H^1(\mathbb{R}^2)$$

where $\gamma : H^1(\mathbb{R}^2) \rightarrow H^{1/2}(\mathbb{R})$ is trace op,

$$\gamma \phi(x_1) = \phi(x_1, 0), \quad x_1 \in \mathbb{R}, \phi \in C_0^\infty(\mathbb{R}^2)$$

Thm (e.g. McLean, 2000, p104)

$$H_{\Gamma}^{-1}(\mathbb{R}^2) = \{\gamma^* \phi \cdot \phi \in H_{\Gamma}^{-1/2}(\mathbb{R})\}$$

where $\gamma^* : H^{-1/2}(\mathbb{R}) \rightarrow H^{-1}(\mathbb{R}^2)$,
defined by γ^* adjoint of γ

$$\langle \gamma^* \phi, \psi \rangle := \langle \phi, \gamma \psi \rangle, \quad \phi \in H^{-1/2}(\mathbb{R}), \psi \in H^{-1}(\mathbb{R}^2)$$

where $\gamma : H^{-1}(\mathbb{R}^2) \rightarrow H^{-1/2}(\mathbb{R})$ is trace op,

$$\gamma \phi(x_1) = \phi(x_1, 0), \quad x_1 \in \mathbb{R}, \phi \in C_0^\infty(\mathbb{R}^2)$$

Given $g \in \dot{H}^{-1}(\mathbb{R}^2)$ find $\phi \in H_{\Gamma}^{-1}$ s.t. $\langle \psi, G\phi \rangle = -\langle \psi, Gg \rangle, \forall \psi \in H_{\Gamma}^{-1}$

$$H_{\Gamma}^{-1}(\mathbb{R}^2) = \{\gamma^* \phi \cdot \phi \in H_{\Gamma}^{-1/2}(\mathbb{R})\}$$

Given $g \in H^{-1}(\mathbb{R}^2)$ find $\phi \in H_{\Gamma}^{-1}$ s.t. $\langle \psi, G\phi \rangle = -\langle \psi, Gg \rangle, \forall \psi \in H_{\Gamma}^{-1}$

$$H_{\Gamma}^{-1}(\mathbb{R}^2) = \{\gamma^* \phi \cdot \phi \in H_{\Gamma}^{-1/2}(\mathbb{R})\}$$

\Leftrightarrow Find $\tilde{\phi} \in H_{\Gamma}^{-1/2}(\mathbb{R})$ s.t.

$$\langle \gamma^* \tilde{\psi}, G\gamma^* \tilde{\phi} \rangle = \langle \gamma^* \tilde{\psi}, Gg \rangle, \quad \tilde{\psi} \in H_{\Gamma}^{-1/2}(\mathbb{R})$$

Given $g \in H^1(\mathbb{R}^2)$ find $\phi \in H_\Gamma^{-1}$ s.t. $\langle \psi, \mathcal{G}\phi \rangle = -\langle \psi, \mathcal{G}g \rangle, \forall \psi \in H_\Gamma^{-1}$

$$H_\Gamma^{-1}(\mathbb{R}^2) = \{\gamma^* \phi \cdot \phi \in H_\Gamma^{-1/2}(\mathbb{R})\}$$

\Leftrightarrow Find $\tilde{\phi} \in H_\Gamma^{-1/2}(\mathbb{R})$ s.t.

$$\langle \gamma^* \tilde{\psi}, \mathcal{G} \gamma^* \tilde{\phi} \rangle = \langle \gamma^* \tilde{\psi}, \mathcal{G}g \rangle, \quad \tilde{\psi} \in H_\Gamma^{-1/2}(\mathbb{R})$$

$$\Leftrightarrow \langle \tilde{\psi}, \gamma \mathcal{G} \gamma^* \tilde{\phi} \rangle = \langle \tilde{\psi}, \gamma \mathcal{G}g \rangle,$$

"

Given $g \in H^{-1}(\mathbb{R}^2)$ find $\phi \in H_{\Gamma}^{-1}$ s.t. $\langle \psi, \gamma \phi \rangle = -\langle \psi, \gamma g \rangle, \forall \psi \in H_{\Gamma}^{-1}$

$$H_{\Gamma}^{-1}(\mathbb{R}^2) = \{ \gamma^* \phi \cdot \phi \in H_{\Gamma}^{-1/2}(\mathbb{R}) \}$$

\Leftrightarrow Find $\tilde{\phi} \in H_{\Gamma}^{-1/2}(\mathbb{R})$ s.t.

$$\langle \gamma^* \tilde{\psi}, \gamma \gamma^* \tilde{\phi} \rangle = \langle \gamma^* \tilde{\psi}, \gamma g \rangle, \quad \tilde{\psi} \in H_{\Gamma}^{-1/2}(\mathbb{R})$$

$$\Leftrightarrow \langle \tilde{\psi}, \gamma \gamma^* \tilde{\phi} \rangle = \langle \tilde{\psi}, \gamma g \rangle,$$

$$S = \gamma \gamma^*$$

Given $g \in H^1(\mathbb{R}^2)$ find $\phi \in H_\Gamma^{-1}$ st $\langle \psi, G\phi \rangle = -\langle \psi, Gg \rangle, \forall \psi \in H_\Gamma^{-1}$

\Leftrightarrow Find $\tilde{\phi} \in H_\Gamma^{-1/2}(\mathbb{R})$ st

$$\langle \tilde{\psi}, S\tilde{\phi} \rangle = \langle \tilde{\psi}, \gamma Gg \rangle, \quad \tilde{\psi} \in H_\Gamma^{-1/2}(\mathbb{R})$$

where

$$S = \gamma G \gamma^* : H^{-1/2}(\mathbb{R}) \rightarrow H^{1/2}(\mathbb{R})$$

Given $g \in \tilde{H}^{-1}(\mathbb{R}^2)$ find $\phi \in \tilde{H}_\Gamma^{-1}$ s.t. $\langle \psi, \mathcal{G}\phi \rangle = -\langle \psi, \mathcal{G}g \rangle, \forall \psi \in \tilde{H}_\Gamma^{-1}$

\Leftrightarrow Find $\tilde{\phi} \in H_\Gamma^{-1/2}(\mathbb{R})$ s.t.

$$\langle \tilde{\psi}, \mathcal{S}\tilde{\phi} \rangle = \langle \tilde{\psi}, \gamma \mathcal{G}g \rangle, \quad \tilde{\psi} \in H_\Gamma^{-1/2}(\mathbb{R})$$

where

$$\mathcal{S} = \gamma \mathcal{G} \gamma^* : H_\Gamma^{-1/2}(\mathbb{R}) \rightarrow H_\Gamma^{1/2}(\mathbb{R})$$

is the (acoustic) single-layer potential operator on $\Gamma_\infty = \{(x_1, 0) \mid x_1 \in \mathbb{R}\} \simeq \mathbb{R}$

Given $g \in \dot{H}^{-1}(\mathbb{R}^2)$ find $\phi \in \dot{H}_\Gamma^{-1}$ st $\langle \psi, G\phi \rangle = -\langle \psi, Gg \rangle, \forall \psi \in \dot{H}_\Gamma^{-1}$

\Leftrightarrow Find $\tilde{\phi} \in \dot{H}_\Gamma^{-1/2}(\mathbb{R})$ st

$$\langle \tilde{\psi}, S\tilde{\phi} \rangle = \langle \tilde{\psi}, \gamma Gg \rangle, \quad \tilde{\psi} \in \dot{H}_\Gamma^{-1/2}(\mathbb{R})$$

where $S = \gamma G \gamma^* : \dot{H}_\Gamma^{-1/2}(\mathbb{R}) \rightarrow \dot{H}_\Gamma^{1/2}(\mathbb{R})$

Given $g \in \tilde{H}^{-1}(\mathbb{R}^2)$ find $\phi \in \tilde{H}_\Gamma^{-1}$ st $\langle \psi, G\phi \rangle = -\langle \psi, Gg \rangle, \forall \psi \in \tilde{H}_\Gamma^{-1}$

\Leftrightarrow Find $\tilde{\phi} \in H_\Gamma^{-1/2}(\mathbb{R})$ st

$$\langle \tilde{\psi}, S\tilde{\phi} \rangle = \langle \tilde{\psi}, \gamma Gg \rangle, \quad \tilde{\psi} \in H_\Gamma^{-1/2}(\mathbb{R})$$

where $S = \gamma G \gamma^* : H_\Gamma^{-1/2}(\mathbb{R}) \rightarrow H_\Gamma^{1/2}(\mathbb{R})$

OK, but still, where are the

INTEGRALS !!

Remember

$$u = Gg$$

$$\Leftrightarrow \Delta u + k^2 u = g$$

Remember

$$u = Gg$$

$$\Leftrightarrow \Delta u + k^2 u = g$$

$$\Leftrightarrow (k^2 - \xi^2) \hat{u}(\xi) = \hat{g}(\xi)$$

$$\Leftrightarrow u = \mathcal{F}^{-1} \left(\frac{\hat{g}(\xi)}{k^2 - \xi^2} \right)$$

Remember $u = Gg$

$$\Leftrightarrow u = \mathcal{F}^{-1} \left(\frac{\hat{g}(\xi)}{k^2 - \xi^2} \right)$$

Remember $u = Gg$

$$\Leftrightarrow u = \mathcal{F}^{-1} \left(\frac{\hat{g}(\xi)}{k^2 - \xi^2} \right)$$

$$\Rightarrow u(x) = \langle g, \Phi(x, \cdot) \rangle$$

if $x \notin \text{supp}(g)$

Hankel fn

or $g \in H^{-1+\epsilon}(\mathbb{R}^2)$

where $\Phi(x) = -\frac{1}{4} H_0^{(1)}(k|x|) \in L^1(\mathbb{R}^2)$

Remember $u = Gg$

$$\Leftrightarrow u = \mathcal{F}^{-1} \left(\frac{\hat{g}(\xi)}{k^2 - \xi^2} \right)$$

$$\Rightarrow u(x) = \langle g, \Phi(x - \cdot) \rangle$$

if $x \notin \text{supp}(g)$

or $g \in H^{-1+\epsilon}(\mathbb{R}^2)$

$$\stackrel{=}{=} \int_{\mathbb{R}^2} g(y) \Phi(x-y) dy$$

$(g \in L^2(\mathbb{R}^2))$

Thus, if $\tilde{\phi} \in L^2(\mathbb{R})$,

$$S\tilde{\phi}(x_1) = (\gamma G \gamma^* \tilde{\phi})(x_1)$$

$$= \int_{\mathbb{R}} \tilde{\Phi}(x_1 - y_1) \tilde{\phi}(y_1) dy_1$$

where $\tilde{\Phi}(t) = -\frac{1}{4} H_0^{(1)}(k|t|), \quad t \in \mathbb{R}$