

The cost of controlling strongly degenerate parabolic equations

Patrick Martinez

Institut de Mathématiques de Toulouse
Université Paul Sabatier, Toulouse III

INDAM - Rome - July 2017 - Piermarco 60

Work in collaboration with

- Piermarco Cannarsa, Univ. Tor Vergata, Roma 2,
- Judith Vancostenoble, Univ. Toulouse 3.

Where the journey has begun...



Outlines

- ① Some examples of degenerate parabolic equations
- ② Controllability cost for a boundary control :
 - upper bounds
 - lower bounds
- ③ Controllability cost for locally distributed control
- ④ Works in progress

Motivation

Properties of null controllability (and inverse problems) of

$$\begin{cases} u_t - \operatorname{div} (A(x) \nabla u) = h(x, t) \chi_\omega, \\ \text{boundary conditions,} \\ \text{initial condition} \end{cases}$$

when Ω bounded domain of \mathbb{R}^n ($n = 1, 2$); ω sub-domain of Ω , and

$$A : \overline{\Omega} \rightarrow M_n(\mathbb{R}), \quad A(x) \text{ symmetric and } \geq 0,$$

but non uniformly positive : for example when

$$\forall x \in \partial\Omega, \quad \det(A(x)) = 0$$

(First step to controllability to trajectories, LQR problems...)

When $A(x)$ is uniformly positive : heat equation **Lebeau-Robbiano (95)**, general case : **Fursikov-Imanuvilov (95, 96)**

Some examples of degenerate parabolic equations in dim 1 :

- climatology : the **Budyko-Sellers** model :

$$RT_t - k((1 - x^2)T_x)_x - S_0s(x)a(x, T) = -I(T), x \in (-1, 1);$$

Ghil (1976...), Diaz (1993...),
Roques-Checkroun-Cristofol-Soubeyrand-Ghil (2014)

- economy : the **Black-Scholes** model :

$$u_t - x^2 u_{xx} + \dots = \dots, x \in (0, L);$$

- combustion theory and quantum mechanics : **inverse square potential**

$$u_t - u_{xx} - \frac{\mu}{x^2} u = 0, x \in (0, 1)$$

Baras-Goldstein (1984), Vásquez-Zuazua (2000),
Vancostenoble-Zuazua (2008).

Some examples in dim 2 :

- ▶ aeronautics : the Crocco equation (boundary layer model) :

$$u_t + a(y)u_x - (b(y)u_y)_y = \text{localized control}, x \in (0, L), y(0, 1)$$

with $a(1) = 0 = b(1)$; Oleinik-Samokhin (1999),
Buchot-Raymond (2002), M-Raymond-Vancostenoble (2003)
(in the simple case where $a(y) = 1$)

- ▶ Kolmogorov type operators :

$$f_t + v f_x - f_{vv} = \text{loc. control}, (x, v) \in (0, 2\pi) \times (0, 2\pi)$$

Beauchard-Zuazua (2009), Beauchard (2014)

- ▶ Grushin type operators :

$$f_t - f_{xx} - x^{2\gamma} f_{yy} = \text{loc. control}, (x, y) \in (-1, 1) \times (0, 1)$$

Beauchard-Cannarsa-Guglielmi (2014)

- An example in biology : the Fleming-Viot model (genetic frequency model) :

$$u_t - \text{Tr} (C(x) D^2 u) \dots = f,$$

where $C(x) = (x_i(\delta_{ij} - x_j))_{i,j}$ and $x \in \{x_i \in [0, 1], \sum_i x_i \leq 1\}$;
example : $N = 2$: $\det C(x) = 0$ along the sides of the triangle.

Cerrai-Clément (2004), Campiti-Rasa (2004),
Albanese-Mangino (2015)

- invariance sets for diffusion processes : Aubin-Da Prato (1990,98) : naturally, the diffusion matrix is degenerate in the normal direction at the boundary (motivation for Cannarsa-M-Vancostenoble, Memoirs AMS (2016)).

The typical strongly degenerate parabolic equation :

Given $\alpha \geq 1$:

- ▶ Locally distributed control :

$$\begin{cases} u_t - (x^\alpha u_x)_x = h(x, t)\chi_{(a,b)}(x), & x \in (0, 1), t > 0, \\ (x^\alpha u_x(x, t))_{/x=0} = 0 = u(1, t) \end{cases},$$

- ▶ Boundary control at the non degeneracy point :

$$\begin{cases} u_t - (x^\alpha u_x)_x = 0, & x \in (0, 1), t > 0, \\ (x^\alpha u_x(x, t))_{/x=0} = 0, & \\ u(1, t) = H(t) \end{cases} :$$

The (theoretical) problem

Known (Cannarsa-M-Vancostenoble (2008)) :

Null Controllability holds if $\alpha \in [1, 2)$, does not hold if $\alpha \geq 2$

Goal : understand the behavior when $\alpha \rightarrow 2^-$.

Natural quantity to estimate : "Null controllability cost" :

$$C(\alpha, T) := \sup_{\|u_0\|=1} \left(\inf_{\text{admissible control : } u(T)=0} \{ \|\text{control}\| \} \right) :$$

$$?? \leq C(\alpha, T) \leq ???$$

expected :

$$C(\alpha, T) \rightarrow +\infty \quad \text{as } \alpha \rightarrow 2^-$$

Related literature

"Controllability cost" appears also in

- ▶ the 'fast control problem' : behavior of $C(T)$ as $T \rightarrow 0$, for several types of equations : Seidman (1984, 2000), Guichal (1985), Miller (2004, 2005, 2006), Tenenbaum-Tucsnak (2007, 2011), Lissy (2014), Benabdallah et al (2014),
- ▶ for semilinear parabolic equations : linearized model (first step) :

$$u_t - \Delta u + a(x, t)u = \dots$$

behavior of $C(\|a\|_\infty, T)$ as $\|a\|_\infty \rightarrow \infty$:
Fernandez-Cara-Zuazua (2000),

- ▶ the 'vanishing viscosity limit' :

$$u_t - \varepsilon u_{xx} + Mu_x = \dots$$

analysis of the importance of the transport term, behavior of $C(\varepsilon, T, M)$ as $\varepsilon \rightarrow 0$, taking care of the size of MT :

Coron-Guerrero (2005), Guerrero-Lebeau (2007), Glass (2010), Lissy (2015),

- ▶ observability cost for 1D wave equation : Haraux-Liard-Privat (2016) ; an equivalent of the observability cost as $T \rightarrow \infty$: Humbert-Privat-Trélat (2016) ;
- ▶ optimizing the location of control region of given measure for 1D wave equation : Privat-Trélat-Zuazua (2013), for 1D heat equation : Privat-Trélat-Zuazua (2017).

2 main complementary methods

2 main methods to study null controllability, and its cost :

► **moment method** :

- Fattorini-Russell (1971, 74)
- eigenvalues, eigenfunctions, biorthogonal families
- **sharp upper and lower bounds** for some equations (1-D, constant coefficients) ;

► **Carleman estimates** :

- Lebeau-Robbiano (1995), Fursikov-Imanuvilov (1996)
- observability
- **good upper bounds** for a large class of equations (N-D, variable coefficients).

Carleman estimates for the degenerate parabolic equation :

$$C(\alpha, T) \leq e^{\frac{C}{(2-\alpha)^4 T}} : \quad \text{bound from below? (sharp estimate?)}$$

Moment method for the boundary control problem

$$\begin{cases} u_t - (x^\alpha u_x)_x = 0, & x \in (0, 1), t > 0, \\ (x^\alpha u_x(x, t))|_{x=0} = 0, & : \\ u(1, t) = H(t) \end{cases}$$

- ▶ well-posedness in weighted Sobolev spaces; $(H \in H^1(0, T))$;
- ▶ eigenvalues, eigenfunctions :

$$\begin{cases} -(x^\alpha \Phi_x)_x = \lambda \Phi, & x \in (0, 1), t > 0, \\ (x^\alpha \Phi_x)|_{x=0} = 0, & : \\ \Phi(1) = 0 \end{cases}$$

then eigenvalues $(\lambda_{\alpha,n})_{n \geq 1}$ associated to $(\Phi_{\alpha,n})_{n \geq 1}$

- ▶ if $u(T) = 0$, then multiplying by $\Phi_{\alpha,n}(x)e^{\lambda_{\alpha,n}(T-t)}$:

$$\forall n \geq 1, \quad \int_0^T H(t)e^{\lambda_{\alpha,n}t} dt = \frac{(u_0, \Phi_{\alpha,n})}{r_{\alpha,n}}, \quad \text{with } r_{\alpha,n} = \Phi'_{\alpha,n}(1);$$

- ▶ if family $(\sigma_{\alpha,m}^+)_{m \geq 1}$ "biorthogonal" to $(e^{\lambda_{\alpha,n}t})_{n \geq 1}$ in $L^2(0, T)$:

$$\forall m, n \geq 1, \quad \int_0^T \sigma_{\alpha,m}^+(t)e^{\lambda_{\alpha,n}t} dt = \delta_{mn} = \begin{cases} 1 & \text{if } m = n, \\ 0 & \text{if } m \neq n. \end{cases}$$

then formally

- ▶ **biorth. family** \implies **control** $H(t) := \sum_{m=1}^{\infty} \frac{(u_0, \Phi_{\alpha,m})}{r_{\alpha,m}} \sigma_{\alpha,m}^+(t)$ drives the solution to 0 in time T ,
- ▶ **control** \implies **biorth. family** : if $u_0 = \Phi_{\alpha,m}$ and H_m is admissible, then $(r_{\alpha,m}H_m)_{m \geq 1}$ is biorthogonal to $(e^{\lambda_{\alpha,n}t})_{n \geq 1}$ in $L^2(0, T)$.

To sum up :

{ upper bound for SOME biorthogonal sequence
information on the eigenfunctions

\implies upper bound for NC cost :

$$C(\alpha, T) \leq \sup_{\|u_0\|=1} \|H\| \leq \left(\sum_{m=1}^{\infty} \frac{\|\sigma_{\alpha,m}^+\|^2}{r_{\alpha,m}^2} \right)^{1/2};$$

{ lower bound for ANY biorthogonal sequence
information on the eigenfunctions

\implies lower bound for NC cost.

Good upper/lower bounds on biorthogonal families?

necessary : information about the eigenvalues and eigenfunctions

Eigenvalues of the degenerate problem

related to Bessel functions and their zeros :

For $\alpha \in [1, 2)$, let

$$\kappa_\alpha := \frac{2 - \alpha}{2}, \quad \nu_\alpha := \frac{\alpha - 1}{2 - \alpha}.$$

Then (Kamke (1948), Everitt-Zettl (1978), Gueye (2014) when $\alpha \in [0, 1)$)

- ▶ the eigenvalues : $\forall n \geq 1, \lambda_{\alpha,n} = \kappa_\alpha^2 j_{\nu_\alpha,n}^2$,
- ▶ the eigenfunctions $\Phi_{\alpha,n}(x) = \frac{\sqrt{2\kappa_\alpha}}{|J'_{\nu_\alpha}(j_{\nu_\alpha,n})|} x^{(1-\alpha)/2} J_{\nu_\alpha}(j_{\nu_\alpha,n} x^{\kappa_\alpha})$,

where

- ▶ J_{ν_α} is the Bessel function of first kind and of order ν_α ,
- ▶ and $(j_{\nu_\alpha,n})_{n \geq 1}$ is the sequence of the positive zeros of J_{ν_α} .

Argument :

λ eigenvalue, Φ associated eigenfunction : then the new function Ψ

$$\Phi(x) =: x^{\frac{1-\alpha}{2}} \Psi\left(\frac{2}{2-\alpha} \sqrt{\lambda} x^{\frac{2-\alpha}{2}}\right)$$

satisfies the following ODE :

$$y^2 \Psi''(y) + y \Psi'(y) + (y^2 - (\frac{\alpha-1}{2-\alpha})^2) \Psi(y) = 0, \quad y \in (0, \frac{2\sqrt{\lambda}}{2-\alpha})$$

which is the Bessel's equation of order ν :

$$y^2 \Psi''(x) + y \Psi'(x) + (y^2 - \nu^2) \Psi(x) = 0, \quad y > 0,$$

with $\nu = \nu_\alpha := \frac{\alpha-1}{2-\alpha}$.

Using the boundary conditions and well-posedness setting, we find $\lambda_{\alpha,n}$ and $\Phi_{\alpha,n}$.

Connection with the other problems :

- ▶ the strongly degenerate parabolic equation :

$$\begin{cases} -(x^\alpha \Phi_x)_x = \lambda \Phi \\ (x^\alpha \Phi_x)(0) = 0 = \Phi(1) \end{cases} \implies \lambda_{\alpha,n} = \kappa_\alpha^2 j_{\nu_\alpha,n}^2,$$

and then

$$\sqrt{\lambda_{\alpha,n+1}} - \sqrt{\lambda_{\alpha,n}} = \kappa_\alpha (j_{\nu_\alpha,n+1} - j_{\nu_\alpha,n}),$$

and (classical for Bessel functions)

$$j_{\nu_\alpha,n+1} - j_{\nu_\alpha,n} \rightarrow \pi \quad \text{as } n \rightarrow \infty,$$

hence

$$\sqrt{\lambda_{\alpha,n+1}} - \sqrt{\lambda_{\alpha,n}} \rightarrow \frac{\pi}{2}(2 - \alpha) \quad \text{as } n \rightarrow \infty.$$

- ▶ the 'vanishing viscosity limit' :

$$\begin{cases} -\varepsilon \Phi_{xx} - M \Phi_x = \lambda \Phi \\ \Phi(0) = 0 = \Phi(1) \end{cases} \implies \lambda_{\varepsilon,n} = \varepsilon \pi^2 n^2 + \frac{M^2}{4\varepsilon} :$$

hence

$$\sqrt{\lambda_{\varepsilon,n+1}} - \sqrt{\lambda_{\varepsilon,n}} \rightarrow \pi\sqrt{\varepsilon} \quad \text{as } n \rightarrow \infty;$$

- ▶ the 'fast control problem' : with the normalization

$$v(x, \tau) := u(x, \tau T) :$$

$$\begin{cases} v_\tau - Tv_{xx} = \text{loc. control}, \\ v_{/x=0} = 0 = v_{/x=1}, \\ v_{/t=0} = u_0, \textcolor{blue}{v_{/t=1} = 0} \end{cases} \implies \textcolor{red}{\lambda_{T,n} = T\pi^2 n^2},$$

and then $\sqrt{\lambda_{T,n+1}} - \sqrt{\lambda_{T,n}} \rightarrow \pi\sqrt{T}$ as $n \rightarrow \infty$.

Existence and bounds of a biorthogonal sequence under 'gap-min' condition

Existence of biorthogonal sequences :

- ▶ Fattorini-Russell (1971, 74) : functional analysis and complex analysis ; gap conditions but not explicit as $T \rightarrow 0$;
- ▶ many applications of their results and methods ;
- ▶ for the dependence $T \rightarrow 0$:
 - Seidman-Avdonin-Ivanov (2000), Tenenbaum-Tucsnak (2007, 2011), Lissy (2014) : complex analysis (but they work with $\lambda_n = rn^2 + l.o.t.$, and do not take care of a large parameter),
 - Glass (2010) : precise, but adaptable to general conditions on the eigenvalues ?

We proved the following version (Cannarsa-M-Vancostenoble (2017)) :

Theorem

Assume that $\lambda_1 \geq 0$, and that

$$\forall n \geq 1, \quad \sqrt{\lambda_{n+1}} - \sqrt{\lambda_n} \geq \gamma_{\min} > 0.$$

Then there exists a family $(\sigma_m^+)_{m \geq 1}$ which is biorthogonal to the family $(e^{\lambda_n t})_{n \geq 1}$ in $L^2(0, T)$, and for which there is some universal constant C_u independent of T , γ_{\min} and m such that

$$\forall m, \quad \|\sigma_m^+\|_{L^2(0, T)}^2 \leq C_u e^{-2\lambda_m T} e^{C_u \frac{\sqrt{\lambda_m}}{\gamma_{\min}}} e^{\frac{C_u}{\gamma_{\min}^2 T}} B(T, \gamma_{\min}),$$

with $B(T, \gamma_{\min}) = \frac{1}{T} \max(T \gamma_{\min}^2, \frac{1}{T \gamma_{\min}^2})$.

Proof : mainly the construction of [Seidman-Avdonin-Ivanov \(2000\)](#) (complex analysis techniques), combined with an additional parameter ([Tenenbaum-Tucsnaak \(2007\)](#), [Lissy \(2015\)](#)) :

Argument

- ▶ a Weierstrass product :

$$F_m(z) := \prod_{k=1, k \neq m}^{\infty} \left(1 - \left(\frac{iz - \lambda_m}{\lambda_k - \lambda_m} \right)^2 \right)$$

whose growth is estimated using the gap condition,

- ▶ a suitable mollifier $M_m(z)$, chosen so that

$$f_m := F_m M_m \text{ satisfies } \begin{cases} \forall m, n \geq 1, & f_m(-i\lambda_n) = \delta_{mn}, \\ \forall z \in \mathbb{C}, & |f_m(-z)e^{-iz\frac{T}{2}}| \leq C_m e^{\frac{T}{2}|z|} \\ \forall m \geq 1, & f_m \in L^2(\mathbb{R}) \end{cases},$$

- ▶ the Paley-Wiener theorem : $f_m(-z)e^{-iz\frac{T}{2}}$ is the inverse Fourier transform of some compactly supported function ϕ_m (properties 2 and 3), that will be the biorthogonal sequence (property 1).

Application : upper bound of the cost

When $\nu_\alpha = \frac{\alpha-1}{2-\alpha} \geq \frac{1}{2}$, the sequence $(j_{\nu_\alpha, n+1} - j_{\nu_\alpha, n})_n$ decays to π ([Komornik-Loreti \(2005\)](#)), hence

$$\sqrt{\lambda_{\alpha, n+1}} - \sqrt{\lambda_{\alpha, n}} \geq \frac{\pi}{2}(2 - \alpha) =: \gamma_{\min}(\alpha),$$

hence the existence of a biorthogonal family, and

$$H^{(u_0)}(t) := \sum_{m=1}^{\infty} \frac{(u_0, \Phi_{\alpha, m})}{r_{\alpha, m}} \sigma_{\alpha, m}^+(t)$$

is well-defined, drives the solution to 0 in time T , hence

$$\begin{aligned} C(\alpha, T) &\leq \sup_{\|u_0\|=1} \|H^{(u_0)}\| \\ &\leq \left(\sum_{m=1}^{\infty} \frac{C_u}{r_{\alpha, m}^2} e^{-2\lambda_{\alpha, m} T} e^{C_u \frac{\sqrt{\lambda_{\alpha, m}}}{\gamma_{\min}(\alpha)}} e^{\frac{C_u}{\gamma_{\min}(\alpha)^2 T}} B(T, \gamma_{\min}) \right)^{1/2} \leq C_u e^{\frac{C_u}{(2-\alpha)^2 T}}. \end{aligned}$$

Lower bound under some global 'gap-max' condition

Motivation :

H_m drives $u_0 := \Phi_{\alpha,m}$ to 0 in time T

$$\implies \text{(moment method)} \int_0^T (r_{\alpha,m} H_m) e^{\lambda_{\alpha,n} t} dt = \delta_{mn},$$

hence $(r_{\alpha,m} H_m)_m$ is biorthogonal to $(e^{\lambda_{\alpha,n} t})_n$.

And bound from below, by hilbertian techniques :

► in \mathbb{R}^2 , \mathbb{R}^3 :

$$(v_1, v_2) \text{ biorthogonal to } (u_1, u_2) \implies \|v_2\| \geq \frac{1}{\text{dist}(u_2, \mathbb{R}u_1)};$$

► similarly in a general context :

- Hansen (1991) : optimality (w.r.t. m) of Fattorini-Russell,
- Guichal (1985) optimality (for heat eq.) of Seidman (1984).

We extend Guichal (1985), the goal being to have

$$\|r_{\alpha,m} H_m\| \geq \dots, \text{ hence } \|H_m\| \geq \frac{\dots}{|r_{\alpha,m}|}, \text{ hence } C(\alpha, T) \geq \frac{\dots}{|r_{\alpha,m}|}.$$

Theorem

(CMV (2017)) Assume that $\lambda_1 \geq 0$, and that there is some $0 < \gamma_{\min} \leq \gamma_{\max}$ such that

$$\forall n \geq 1, \quad \gamma_{\min} \leq \sqrt{\lambda_{n+1}} - \sqrt{\lambda_n} \leq \gamma_{\max}.$$

Then there exists some $C(m, \gamma_{\max}, \lambda_1) > 0$ (given explicitly in the paper) and $c_u > 0$ independent of T and m such that : **any** family $(\sigma_m^+)_{m \geq 1}$ which is biorthogonal to the family $(e^{\lambda_n t})_{n \geq 1}$ in $L^2(0, T)$ satisfies :

$$\|\sigma_m^+\|_{L^2(0, T)}^2 \geq e^{-2\lambda_m T} e^{\frac{1}{2\gamma_{\max}^2 T}} b(T, \gamma_{\max}, m),$$

with

$$b(T, \gamma_{\max}, m) = \frac{c_u^2}{C(m, \gamma_{\max}, \lambda_1)^2 T} \left(\frac{1}{2\gamma_{\max}^2 T} \right)^{2m} \frac{1}{(4\gamma_{\max}^2 T + 1)^2}.$$

Argument

- ▶ As in \mathbb{R}^2 : **any** biorthogonal sequence verifies

$$\|\sigma_m^+\| \geq \frac{1}{d_{T,m}}, \quad \text{where } d_{T,m} := \text{dist} (e^{\lambda_m t}, \overline{\text{Vect} \{e^{\lambda_k t}, k \neq m\}});$$

- ▶ then Guichal (1985) :

$$d_{T,m} \leq \|e^{\lambda_m t} - \sum_{i=1, i \neq m}^{M+1} A_i e^{\lambda_i t}\| = \|\frac{-1}{\tilde{A}_m} \sum_{i=1}^{M+1} -\tilde{A}_i e^{\lambda_i t}\| = \|\frac{-1}{\tilde{A}_m} q(t)\|,$$

with a special choice of the coefficients \tilde{A}_i : chosen such that

$$q(0) = \dots = q^{(M-1)}(0) = 0, \quad q^{(M)}(0) = 1$$

(in order to have q small) ;

- ▶ then (Vandermonde determinant)

$$d_{T,m} \leq \left(\prod_{i=1, i \neq m}^{M+1} |\lambda_i - \lambda_m| \right) \left(\int_0^T \frac{s^{2M}}{M!^2} e^{-2\lambda_1 s} ds \right)^{1/2} \leq \dots$$

Application

As already seen :

$$\pi \leq j_{\nu_\alpha, n+1} - j_{\nu_\alpha, n} \leq j_{\nu_\alpha, 2} - j_{\nu_\alpha, 1},$$

hence $\gamma_{\max} = \kappa_\alpha (j_{\nu_\alpha, 2} - j_{\nu_\alpha, 1})$. But (classical)

$$j_{\nu, 2} - j_{\nu, 1} \sim a\nu^{1/3} \quad \text{as } \nu \rightarrow +\infty.$$

Hence $\gamma_{\max} \sim c\nu_\alpha^{-2/3} \sim c'(2-\alpha)^{2/3}$ as $\alpha \rightarrow 2^-$, hence

$$C(\alpha, T) \geq \|H_1(\text{ driving } \Phi_{\alpha, 1})\| \geq \frac{\dots}{|r_{\alpha, 1}|} e^{\frac{1}{2\gamma_{\max}^2} T} = \frac{\dots}{|r_{\alpha, 1}|} e^{\frac{c}{(2-\alpha)^{4/3}} T};$$

hence

$$e^{\frac{c}{(2-\alpha)^{4/3}} T} \leq C(\alpha, T) \leq e^{\frac{c}{(2-\alpha)^2} T}.$$

Hence $C(\alpha, T) \rightarrow \infty$ as $\alpha \rightarrow 2^-$, BUT gap between $\frac{4}{3}$ and 2 ...

Additionnal property of the zeros of Bessel functions

The gap $(j_{\nu_\alpha, n+1} - j_{\nu_\alpha, n})_{n \geq 1}$ decays

- ▶ from $j_{\nu_\alpha, 2} - j_{\nu_\alpha, 1} \sim a\nu_\alpha^{1/3}$ (large)
- ▶ to π .

Hence certainly :

$$\exists N_\nu, \forall n \geq N_\nu, \quad j_{\nu_\alpha, n+1} - j_{\nu_\alpha, n} \leq 2\pi,$$

but this was not taken into account. The value of N_ν :

Lemma

(CMV (2017))

$$\forall \nu \geq \frac{1}{2}, \forall n > \nu, \quad j_{\nu_\alpha, n+1} - j_{\nu_\alpha, n} \leq 2\pi.$$

Proof : oscillation theorem of Sturm for second order ODE, in the spirit of Komornik-Loreti (2005).

Lower bound under some asymptotic 'gap-max' condition

Theorem

Assume that $\lambda_1 \geq 0$, and that there are $0 < \gamma_{\min} \leq \gamma_{\max}^* \leq \gamma_{\max}$ such that

$$\forall n \geq 1, \quad \gamma_{\min} \leq \sqrt{\lambda_{n+1}} - \sqrt{\lambda_n} \leq \gamma_{\max},$$

and

$$\forall n \geq N_*, \quad \sqrt{\lambda_{n+1}} - \sqrt{\lambda_n} \leq \gamma_{\max}^*.$$

Then **any** family $(\sigma_m^+)_{m \geq 1}$ which is biorthogonal to the family $(e^{\lambda_n t})_{n \geq 1}$ in $L^2(0, T)$ satisfies :

$$\|\sigma_m^+\|_{L^2(0, T)}^2 \geq e^{-2\lambda_m T} e^{\frac{2}{T(\gamma_{\max}^*)^2}} b^*(T, \gamma_{\max}, \gamma_{\max}^*, N_*, \lambda_1, m)^2$$

(where b^* is explicitly given in our paper).

Application

Applying the previous result with the asymptotic gap of the zeros of Bessel functions :

- ▶ $\gamma_{max} = j_{\nu_\alpha,2} - j_{\nu_\alpha,1} \sim a\nu_\alpha^{1/3},$
- ▶ $\gamma_{max}^* = 2\pi,$
- ▶ $N_* = [\nu_\alpha] + 1,$

we obtain

$$e^{-\frac{1}{c} \frac{1}{(2-\alpha)^{4/3}} (\ln \frac{1}{2-\alpha} + \ln \frac{1}{T})} e^{\frac{c}{T(2-\alpha)^2}} \leq C(\alpha, T) \leq e^{\frac{c}{T(2-\alpha)^2}}.$$

The locally distributed control problem

$$\begin{cases} u_t - (x^\alpha u_x)_x = h(x, t) \chi_{(a,b)}(x), & x \in (0, 1), t > 0, \\ (x^\alpha u_x(x, t))|_{x=0} = 0 = u(1, t) \end{cases}$$

Similar :

- ▶ NC holds iff $\alpha < 2$;
- ▶ formally admissible control :

$$h(x, t) = h(x, t) := \sum_{m \geq 1} -(u_0, \Phi_{\alpha, m}) \sigma_{\alpha, m}^+(t) \frac{\Phi_{\alpha, m}(x)}{\int_a^b \Phi_{\alpha, m}^2} ;$$

- ▶ hence information needed for upper estimate of the NC cost :
lower bound of $\int_a^b \Phi_{\alpha, m}^2$;
- ▶ classical (Lagnese (1983)) : $\inf_m \int_a^b \Phi_{\alpha, m}^2 > 0$, but not sufficient to estimate $\|h\|$ in function of α ;

- ▶ we prove the following :

Proposition

There exists $\gamma_0^* > 0$ independent of $\alpha \in [1, 2)$ and of $m \geq 1$ s.t.

$$\forall \alpha \in [1, 2), \forall m \geq 1, \quad \int_a^b \Phi_{\alpha, m}^2 \geq \gamma_0^*(2 - \alpha);$$

Difficulties

- in our case

$$\int_a^b \Phi_{\alpha,m}^2 = \frac{2}{j_{\nu_\alpha,m}^2 |J'_{\nu_\alpha}(j_{\nu_\alpha,m})|^2} \int_{j_{\nu_\alpha,m}}^{j_{\nu_\alpha,m} b^{\kappa_\alpha}} y J_{\nu_\alpha}(y)^2 dy,$$

- known : $J_\nu(y)$ either as $y \rightarrow +\infty$ or as $\nu \rightarrow +\infty$,
- but here y and ν go to $+\infty$ simultaneously ;

- ▶ and proof :
 - the Bessel ODE,
 - integral representation of solutions (keeping in mind ν_α large)
 - classical properties of sin (Haraux (1978))
- ▶ and then similar upper and lower estimates for the NC cost.

So to conclude :

- ▶ many properties of Bessel functions to use,
- ▶ some of them to complete,
- ▶ and explicit and quite practical results for biorthogonal sequences under global/asymptotic gap conditions.

(The reaction of my son, a few years ago, when he opened the Watson's book :)




Figure – (Bonne fête, Florent !)

Works in progress and open questions

- ▶ **degeneracy inside the domain** : NC and its cost in function of the degeneracy parameter (joint work with P. Cannarsa and Roberto Ferretti)
- ▶ **inverse square potential** : precise estimates of the cost in function of the parameter μ (joint work with J. Vancostenoble)
- ▶ some of this useful to prove the optimality of the minimal time for the **Grushin operator** ??
- ▶ (more difficult) pick a number n between 1 and 10, and try to guess in **which country (continent!)** Piermarco will be in n weeks ;-)

Works in progress and open questions

- ▶ **degeneracy inside the domain** : NC and its cost in function of the degeneracy parameter (joint work with P. Cannarsa and Roberto Ferretti)
- ▶ **inverse square potential** : precise estimates of the cost in function of the parameter μ (joint work with J. Vancostenoble)
- ▶ some of this useful to prove the optimality of the minimal time for the **Grushin operator** ??
- ▶ (more difficult) pick a number n between 1 and 10, and try to guess in **which country (continent!)** Piermarco will be in n weeks ;-)

Works in progress and open questions

- ▶ **degeneracy inside the domain** : NC and its cost in function of the degeneracy parameter (joint work with P. Cannarsa and Roberto Ferretti)
- ▶ **inverse square potential** : precise estimates of the cost in function of the parameter μ (joint work with J. Vancostenoble)
- ▶ some of this useful to prove the optimality of the minimal time for the **Grushin operator** ??
- ▶ **(more difficult)** pick a number n between 1 and 10, and try to guess in **which country (continent!)** Piermarco will be in n weeks ;-)

Thank you for your attention, and most of all

Happy birthday, Piermarco !